

# Annual Ryegrass Control in Hayfields

*Dr. Patrick McCullough, Extension Specialist, the University of Georgia*

Annual ryegrass (*Lolium multiflorum*), also referred to as Italian ryegrass, is the most problematic winter annual weed in Georgia hayfields. Seed germinates from September to November when soil temperatures drop below 70° F. Seedlings mature in fall, overwinter in a vegetative state, and resume active growth in spring. Annual ryegrass grows well under cool conditions when pasture grasses are dormant or have limited competitive growth. Plants exhibit erect growth that reaches approximately three feet in height upon maturity (Picture 1, 2). Consequently, annual ryegrass may interfere with pasture grasses during spring green-up and beyond. Annual ryegrass also may absorb potassium (K) and other crucial plant nutrients during the spring. Such nutrient absorption can negatively influence the summer forage's production, as this may induce nutrient deficiencies in the pasture grass long after the annual ryegrass has died. There is also significant evidence that stressed annual ryegrass (e.g., drought stress, late-season disease pressure, etc.) may cause it to release allelopathic compounds (natural plant chemicals that inhibit the growth of other plants in their vicinity) that are active against bermudagrass and other desirable forages. Research in North Carolina has indicated that this interference from ryegrass may reduce the yield of hayfields by as much as 50%. Furthermore, areas with heavy infestations of annual ryegrass are predisposed to invasion by summer annual weeds that may further exacerbate the yield reduction and accelerate the decline of hayfield stands.

Annual ryegrass is a prolific seed producer that contributes to annual infestations. The seedhead is a long spike with at least 10 alternating florets (Picture 3). The florets contain long awns that are not present on perennial ryegrass (*Lolium perenne*). Plants typically produce seedheads in March in most parts of Georgia. Seed dispersed in late spring can remain dormant in soil for years. Another key characteristic to help identify annual ryegrass, without seedheads, is the clasping auricle at the junction of the leaf sheath and blade (Picture 4). The auricle may help growers identify an annual ryegrass from other weedy grasses, such as tall fescue.

### Cultural Control

Growers can make modifications to their management programs to reduce annual ryegrass establishment in fall. For example, nitrogen fertilization should be reduced during peak germination and periods of vigorous growth. High nitrogen rates in winter and spring will encourage annual ryegrass seed production, dispersal and survival. Mowing is an important cultural practice that can reduce the competitive growth of problematic weeds with hayfield grasses. Mowing before seedhead formation can suppress annual ryegrass growth and

inhibit the production of viable seed in spring. However, the costs and benefits of mowing should be weighed against the costs and benefits of preventive or curative chemical applications.

Practices that disrupt soil – such as aeration, sub-soiling, or tilling operations – should be conducted when tall fescue, bermudagrass, or bahiagrass stands are actively growing. Voids left in fields with exposed soil may permit annual ryegrass invasion during periods of peak germination. Timing these operations during favorable periods for quick recovery promotes competition with weeds. In tall fescue hayfields, growers can promote recovery from summer stresses before annual ryegrass germination in fall. Reseeding tall fescue in thinned areas to promote a dense stand promotes competition with annual ryegrass seedlings in fall to reduce infestations.

### Chemical Control

Growers may use Prowl H<sub>2</sub>O 3.8SC (pendimethalin) in established bermudagrass, bahiagrass, alfalfa, and cool-season forage grasses, including tall fescue, for pre-emergence control of annual ryegrass in hayfields (Table 1). Applications should be timed when soil temperatures drop below 70° in fall. Prowl may be applied at 2 to 3 qt per acre for annual ryegrass control. Treatments should be applied prior to a rainfall to enhance soil incorporation and herbicide activation. Prowl is also not labeled for perennial peanut or forage legumes, other than alfalfa, grown for hay. Sequential treatments may be applied at a 6- to 8-week interval in fall to extend the length of control in labeled species.

The optimum timing for post-emergence control of annual ryegrass is when plants are less than 6 inches in height in fall. Bermudagrass growers may use Impose 2AS (imazapic) or Pastora 71.2DF (nicosulfuron + metsulfuron) for annual ryegrass control, but these products are not labeled for bahiagrass and tall fescue due to excessive injury potential (Table 1). Post-emergence herbicides are more effective in early winter, compared with spring timings, because of the size and maturity of plants at application. Annual ryegrass is generally susceptible to post-emergence herbicides in early winter prior to the onset of freezing temperatures and before seedhead emergence.

Pastora is more efficacious than Impose at labeled use rates in bermudagrass hayfields. Pastora 71.2DF at 1 to 1.25 oz/acre is recommended in November or early December in bermudagrass. Impose 2AS (imazapic) controls annual ryegrass in fall, but requires high rates (8 to 10 oz/acre) on plants greater than 6 inches in height. Bermudagrass is also more susceptible to injury from Impose than Pastora, especially at rates required

# Expert Advice

Table 1. Herbicides for annual ryegrass control in hayfields.

Application Timing	WSSA Group <sup>a</sup>	Herbicide	Labeled Species	Labeled Rate	Control <sup>b</sup>
Pre-emergence	3	pendimethalin (Prowl H <sub>2</sub> O 3.8SC)	Alfalfa	3.1 to 4.2 qt/acre	G
			Bahiagrass		
			Bermudagrass		
Post-emergence	2	imazapic (Impose 2AS)	Bermudagrass	6 to 10 oz/acre	P-F
		nicosulfuron + metsulfuron (Pastora 71.2DF)	Bermudagrass		
	9	glyphosate (Roundup, Accord, others)	Bermudagrass	see label	G-E

<sup>a</sup>WSSA group numbers: 2 = acetolactate synthase (ALS) inhibitor, 3 = mitotic inhibitor, 9 = EPSP synthase inhibitor.

<sup>b</sup>Efficacy for control ratings: P = poor (less than 70%), F = fair (70 to 79%), G = good (80 to 89%), E = excellent (90 to 100%).



**Picture 1:**

Annual ryegrass clumps in a field in spring.



**Picture 2:**

Annual ryegrass plant.



**Picture 3:**

Seedheads of annual ryegrass.



**Picture 4:**

The auricle of annual ryegrass.

for annual ryegrass control. Both herbicides require a non-ionic surfactant at 0.25% v/v of spray solution (1 qt/100 gal).

If growers must apply herbicides in January or February, it is recommended that Pastora treatments have a tank-mixture with glyphosate to enhance control. Moderate rates of glyphosate in bermudagrass, such as 0.125 to 0.25 lb ae/acre, generally do not affect spring transition when applied in winter. However, glyphosate use in spring could cause delayed green-up and growth inhibition to bermudagrass. Sethoxydim (Poast, others) controls ryegrass in dormant bermudagrass, but is not labeled for hayfields.

Growers have no selective herbicides available for post-emergence control of annual ryegrass in bahiagrass or tall fescue. Spot treatments of glyphosate are recommended in these species for control. A 0.5% to 1% solution of glyphosate should be sprayed to annual ryegrass shoots until runoff occurs. Growers should check the product label of certain glyphosate products, and include an adjuvant at 1% v/v if needed.

### Herbicide Resistance

A major limitation to post-emergence control of annual ryegrass is herbicide resistance. Resistance of annual ryegrass

to acetolactate synthase inhibitors (i.e. Impose, Pastora), acetyl CoA-carboxylase inhibitors (sethoxydim, others), and glyphosate has been confirmed in Georgia. Resistance develops from selection pressure by repeated use of the same herbicide or mode of action over years. Many biotypes of annual ryegrass may be present in a population. Genetic differences among biotypes contribute to susceptibility levels to herbicides through altered target-site binding or enhanced degradation. Other resistance mechanisms for annual ryegrass may include reduced absorption, herbicide sequestration, or overproduction of the target-site enzyme.

As susceptible biotypes are controlled by a particular herbicide over years, resistant biotypes may spread in these fields. This type of selection pressure will shift an annual ryegrass population from susceptible to resistant biotypes over years. Growers should have an appreciation for the potential development of resistance to Pastora and glyphosate in annual ryegrass populations. Pre-emergence control of annual ryegrass with Prowl will help delay resistance to herbicides available in hayfields. 