

Methods for Vegetative Establishment



Dr. Dennis Hancock

Extension Forage Specialist

Crop and Soil Sciences – UGA

Most Common Establishment Mistakes



1. Failure to soil test, lime, & fertilize

Most Common Establishment Mistakes

2. Poor seedbed preparations

- "Prepared" seedbed (conv. tillage)
- Sod destruction (no till; minimum till)

Most Common Establishment Mistakes



3. Plant too deep or too shallow

Most Common Establishment Mistakes

4. Poor pest control

Most Common Establishment Mistakes

5. Wrong time of year

Most Common Establishment Mistakes

6. Poor quality seed or vegetative material


Most Common Establishment Mistakes



7. Uncooperative weather

A little perspective...


Sprigging stick designed to push bermudagrass sprigs into the ground.



Two-row sprig planter developed at the CPES by UGA's James Stephens.

Dr. Glen Burton

"Father" of forage & turf bermudagrasses
(1910-2005)



ESTABLISHING THE TIFTON HYBRID BERMUDAGRASSES

Glenn W. Burton

For much of the South, the Tifton hybrid bermudagrasses top the list of forages that may be grown for hay or grazing. Properly managed, they are more dependable and produce more hay or animal product per acre at a lower cost per ton of hay or pound of beef or milk than any forage I know. They contain more dry matter when cut for hay (25 to 30%) and cure faster than other forages. Planted and managed properly, the hybrid bermudagrasses can become well established and provide grazing or hay in the first season. A description of steps required for success follow:



1: Choose an appropriate site for establishment.



- The soil must be well-drained.
- Choose site that is as weed-free as possible.
- Preferably, site is free of bermudagrass or bahiagrass.
- If either are present:
 - Grow a summer crop for 1-2 seasons
 - Chemical fallow (non-selective herbicide 2-4 times during season)
 - High rate of glyphosate in fall prior to spring establishment.



If Bermudagrass or Bahiagrass is Present...



- Use a smother crop for at least one year.
 - Pearl millet, sorghum x sudan
 - RR soybeans
- If replacing a bermudagrass with a different bermudagrass variety, consider two years.



2: Soil test and lime and fertilize accordingly.



- Ask for recommendations for “hybrid bermudagrass hayfield” even if for a pasture.
 - Lime to a target of pH 6.5.
 - Incorporate all recommended nutrients.
- Soil pH of 6.0-7.0 in winter before planting or not a viable field.
- Immediately prior to planting:
 - 50 lbs N, 15 lbs P₂O₅, and 100 lbs K₂O/a
 - Incorporate (no deeper than 2 inches)



Conventional Seedbed Preparation Steps

4. Plow/disc/finish at least 4 wks prior to planting or sprigging
5. Incorporate phosphorus, potassium, and additional lime (as recommended by soil test).
6. Allow time to settle or firm with cultipacker/roller.



Seedbed should be firm

- Boot tracks should be ~1/4 in. deep
- If too fluffy, the soil will dry very quickly (sandy soils)





3: Use the false or stale seedbed preparation method.




- In the 4-6 weeks between tillage and planting, weed seeds will germinate.

False seedbed prep:

- Kill the weeds by lightly tilling the soil with a light drag or shallow disking, then immediately firm with a roller.
 - Downside: moisture loss

Stale seedbed prep:

- Kill the weeds with non-selective herbicide, wait 1-7 days and plant.



Minimum Till (“No-till”) Options

- Acceptable for vegetative establishment (sprigging), but will leave the field very rough.
- Weed control?




4: Choose an establishment method and timing.



Three primary methods

- 1) Dormant sprigs – 40-70 bu of LS/acre
 - Jan. to early Mar.
 - Less desirable for Tifton 85
 - Cover with at least 2" of soil to protect sprigs from freezing
 - 50%+ of dormant sprigs fail to emerge
 - Fall prior: do not allow the nursery area to be cut or grazed after Labor Day
 - Excessive winter rainfall limits dormant sprig survival
 - Estimate sprig survival by grow-out and adjust sprigging rate accordingly



4: Choose an establishment method and timing.



Three primary methods

2) Spring sprigs – 40-70 bu of LS/acre

- Spring (after last freeze) to early Aug.
- Early sprigging increases likelihood of establishment by end of the first year
- Avoid planting before early April
- Sprigs should be vigorously growing before digging.
- Stand development is directly proportional to sprigging rate









4: Choose an establishment method and timing.



Three primary methods

3) Tops/green stems – 60-100 bu/A

- June until early Aug.
- Tops need 6+ nodes on the stolons
 - Fine-textured varieties: 10-12"
 - Coarse-textured varieties: 18-24"
- Nursery area should receive: 100 lbs N, 25 lbs P_2O_5 , and 100 lbs of K_2O /acre in late March to produce tops by June
- Not recommended for Tifton 44
- Usually not planted with sprig planter.





Photo credit: Dr. Yoana Newman, Univ. of Florida



Photo credit: Dr. Yoana Newman, Univ. of Florida



Photo credit: Dr. Yoana Newman, Univ. of Florida





Photo credit: Dr. Yoana Newman, Univ. of Florida



Photo credit: Dr. Yoana Newman, Univ. of Florida



Photo credit: AGCO Corp.



Photo credit: Dr. Yoana Newman, Univ. of Florida



Photo credit: Dr. Yoana Newman, Univ. of Florida

5: Plant ONLY in moist soil.



- Sprigs will die if they drop below **~50-55% moisture** or if they heat above **120°F** for extended period.
- If soil is dry, especially if hot, it will draw moisture out of the sprigs even after they have been planted.
- Ideal: planting on cool, cloudy day, preferably with a misty rain or imminent rainfall.
- Irrigation before and after can add flexibility, but do not over irrigate.
 - ~1"/wk (0.5" x 2x/wk) for first 4 wks



6: Plant pure sprigs or tops.



- Recommended to buy only certified planting material
 - GA Crop Improvement Assoc. certified (www.georgiacrop.com)
- If none available in your area, ask to see the nursery field
 - 'Common' contamination is common complaint.
 - Be proactive!




7: Plant fresh sprigs or tops from a well-fertilized nursery.



Recommended nursery protocol:

- If dormant sprigs to be harvested
 - 100 lbs N, 25 lbs P_2O_5 , and 100 lbs K_2O /acre in Sept. prior to dig.
- If spring sprigs to be harvested
 - 100 lbs N, 25 lbs P_2O_5 , and 100 lbs K_2O /acre at spring green up or within 6 weeks of digging or top harvest.



What Does a “Good Sprig” Look Like?

- Crown, rhizome, and stolon size are an indicator of CHO storage and vigor.
- Crown and rhizomes should be $\frac{3}{16}$ – $\frac{3}{8}$ ” in diameter.
- Stolons in sprigs should have 2-4+ nodes and be $\frac{1}{8}$ – $\frac{1}{4}$ ”+ in diameter at nodes.
- Stolons as tops should have 6+ nodes and be $\frac{1}{8}$ – $\frac{1}{4}$ ”+ in diameter at nodes.
- Dormant and spring sprigs should be planted ASAP (at least <24 hr after digging).
- Tops should be planted < 4 hr of cutting.



8: Pack the soil well after planting.







9: Spray to control weeds.

- Aggressive sprigs planted at high rate can usually out-compete weeds
- If for turf or roadside, more flexible herbicide options
- Mowing (when 7-8" tall, mow to 3")
- Diuron (e.g., Direx 4L @ 1-2.4 qt/a)
 - Good control: crabgrass, crowfootgrass, sandbur, goosegrass, and some broadleaf weeds
 - Immediately after planting
 - Sprigs must be 2-3" deep
 - Tifton 85 is very sensitive to diuron
- 2,4-D (2 qts/a) or 2,4-D + dicamba (2-4 pts/a)
 - Good control of broadleaf weeds
 - No demonstrable pre-emergent activity





10: Complete keys 5-9 on the same day.

- Ensures adequate soil moisture
- Maximizes sprig survival
- Minimizes weed problems



What is a Good Stand of Sprigged Bermudagrass at 1 MAP?

- **At 1 month after planting:**
 - **Healthy plant (4" diameter of growth) w/in a natural stride from one another (< 2.5'), &**
 - **Stolons extending 6-12"+ from most plants**
 - **This equates to roughly 20% stand**
- **If this is the case, apply 50-60 lbs N/a every 4-6 wk & 100 lbs K₂O/a in late July.**
- **Stimulate stolon growth by mowing regularly (7" -> 3") until early Sept.**

What is a Good Stand of Sprigged Bermudagrass in August?

- **ROT: If planted in June (weather permitting), sprigged stands should have roughly the same density as a % as the bu/acre sprigging rate by middle of Aug.:**
 - **Planted June 10 @ 66 bu/a -> 66% stand**
 - **Planted June 15 @ 50 bu/a -> ~50% stand**

Resources



ESTABLISHING BERMUDAGRASS FROM SPRIGS OR TOPS

*Dennis W. Hancock, PhD
Forage Extension Specialist,
Crop and Soil Sciences Department*

Over the last couple of years, I have received many questions about how to establish bermudagrass. Hybrid bermudagrass varieties produce 20-50% more forage than seeded bermudagrasses. So, many producers seek to establish the hybrids. However, hybrid bermudagrasses produce few viable seed and must be established from vegetative plant material (e.g., sprigs or tops). Significant acreage in Georgia was planted to bermudagrass in 2015, and I suspect more will be planted this year, as well. So, how exactly is bermudagrass vegetatively established?



THE UNIVERSITY OF GEORGIA
COLLEGE OF AGRICULTURAL & ENVIRONMENTAL SCIENCES

STUDENTS ALUMNI DEPARTMENTS EXTENSION RESEARCH PUBLICATIONS TOPICS A-Z CALENDAR NEWS

- ∨ CAES Home
- ∨ Commodities
- ∨ Field Crops
- ∨ **Forages**
 - Adapted Species
 - Establishment Guidelines
 - Fertilization Guidelines
 - Pest Management
 - Publications
 - Popular Press Archive
 - Forage Team
 - **FAQs**
 - Related Links

[Commodities](#): [Field Crops](#): [Forages](#): [FAQs](#)

Sprigging Bermudagrass

Who's available to sprig bermudagrass in Georgia?

This webpage provides a list of people, farms, or companies that provide sprigs and/or sprigging services. Information herein was provided by the persons listed or their County Extension Agent. If a section is blank, the submission for that section was blank. The listing of these persons or entities on this page does not imply endorsement by UGA Extension, Dr. Hancock, or the University of Georgia. This is merely a listing provided to help those who need sprigs or sprigging services connect with those in their area who can perform that service. If you would like to be added to this list, please call (706-310-3464) or [email](#) Cathy Felton in our office from 8 a.m. to 12 p.m.

Counties

[Brantley](#), [Burke](#), [Chattooga](#), [Colquitt](#), [Crisp](#), [Effingham](#), [Greene](#), [Jefferson](#), [Lamar](#), [Miller](#), [Morgan](#), [Muscogee](#), [Pulaski](#), [Screven](#), [Thomas](#), [Upson](#), [Wheeler](#)

Questions?



www.georgiaforages.com