


2019 Baleage and Silage Short Course:
The Uptake, Mode of Action, and Fate of
Herbicides in Hayfields

Dr. Tim Gray
UGA Prof. and Research Weed Scientist




The Uptake, Mode of Action, and Fate of Herbicides Used in Hayfields
T.L. Grey
University of Georgia

Mode of Action - Terminology

- **Mode of Action:**
 - How a particular herbicide acts on a plant
 - Response of plant to phytotoxic effects of the herbicide
 - How the plant responds to the herbicide

Plant target Sites



Labels include: Golgi vesicles, ribosome, smooth ER (no ribosomes), nucleolus, nucleus, rough ER (endoplasmic reticulum), large central vacuole, amyloplast (starch grain), cell wall, cell membrane, Golgi apparatus, chloroplast, vacuole membrane, raphide crystal, druse crystal, mitochondrion, cytoplasm.

Mode of Action

- **Primary Mechanism of Action:** plant processes affected by lowest phytotoxic dose of herbicide.
- **Secondary Mechanism of Action:** other plant processes affected by herbicide.

- **Description of MOA**
- <http://www.wssa.net/Weeds/Resistance/WSSA-Mechanism-of-Action.pdf>

Summary of Herbicide Mechanism of Action According to the Weed Science Society of America (WSSA)

1 Acetyl CoA Carboxylase (ACCase) inhibitors
Aryloxyphenoxypyrazone (PQP), cyclohexanedione (DHQ) and phenylpyrazolin (DPA) herbicides inhibit the enzyme acetyl-CoA carboxylase (ACCase), the enzyme catalyzing the first committed step in de novo fatty acid synthesis (Burton 1989; Focke and Lichtenhaler 1987). Inhibition of fatty acid synthesis presumably blocks the production of phospholipids used in building new membranes required for cell growth. Broadleaf species are naturally resistant to cyclohexanedione and aryloxyphenoxypyrazone herbicides because of an insensitive ACCase enzyme. Similarly, natural tolerance of some grasses appears to be due to a less sensitive ACCase (Stoberberg 1989). An alternative mechanism of action has been proposed involving destruction of the electrochemical potential of the cell membrane, but the contribution of this hypothesis remains in question.

2 Acetolactate Synthase (ALS) or Acetylhydroxy Acid Synthase (AHAS) inhibitors

Herbicide Mode of Action - WSSA


- Group herbicides by plant processes affected:
 - Acetyl CoA Carboxylase Inhibitors (1) – **sethoxydim**
 - Amino acid synthesis inhibitors (2) – **SU's, IMI's**
 - Microtubule assembly inhibition (3) - **pendimethalin**
 - Photosynthetic inhibitors (5, 6, 7, 22) – **diuron, paraquat, metribuzin**
 - EPSP synthesis (9) - **glyphosate**
 - PPO (14) - **flumioxazin**



Question

- What happens to herbicides?
- How do these and other herbicides dissipate when applied?
- Limited information in forages
- We know the properties

Environmental fate



1 – ACCase

Acetyl CoA Carboxylase Inhibitors

- Grass herbicides that we use in legumes many times
- Inhibit lipid production
- POST applied
- FOPS & DIMS
- sethoxydim
- Poast
- Resistance issues!!!
- No to low residual



2 - Amino acid inhibitors

- essential building blocks for plant growth and function
- unlike animals, plants make their own
- amino acids are the primary components of proteins and nucleic acids
- proteins are generally storage proteins or enzymes

2 - Amino Acid Inhibitors

- generally target a specific enzyme
 - Some block vital steps in the formation of amino acids- proteins, enzymes
 - branched chain amino acid inhibitors
 - Leucine, Isoleucine, Valine
- dependent on plant growth for activity
 - better growth - better control, slow death
- systemic herbicides
- Soil activity
 - None (imazamox)
 - Some (Metsulfuron, nicosulfuron)
 - Long activity – (imazapic)

Imazethapyr - imazapic



johnsongrass



Ivyleaf morningglory

ALS Inhibitors

- **Imidazolinones**
 - Imazapic (Impose)
 - Imazamox (Raptor)
 - Imzethapyr (Pursuit)
- **Sulfonylureas**
 - sulfosulfuron
 - Nicosulfuron + metsulfuron (Pastora)
 - Many others

Organic matter & clay

- **Positive correlation between sorption & organic matter content**
 - † OM increase, sorption increase
- **Alkaline soils with low OM**
 - † SU degrade slowly
- **Sulfosulfuron, chlorsulfuron reported**
- **Clay mineral sorption – varies from none to some**

Leaching

- **SU herbicides can be mobile in soil**
 - † Experiments have demonstrated
- **R_f values from 0.21 to 0.9**
 - † Chlorsulfuron
 - † Metsulfuron
 - † Sulfometuron
- **Primarily dependent on soil type & characteristics – pH, OM, etc.**
- **Never been a major concern low rates**


SU facts

- **Most all are formulated as WP or DG**
- **Photolysis minor**
- **Volatilization minor**
- **Can move upward even when they were not previously detected**
 - via capillary soil water flow

Conclusions

- **↑ soil pH ↑ SU persistence**
- **↑ temperature ↑ soil dissipation**
- **↑ soil OM content ↓ plant availability**
- **Low use rates combined with factors above**
- **Low leaching potential**

9 – EPSP synthase Glyphosate



- **broadspectrum postemergence weed control**
- **glyphosate labeled in multitude of areas**
- **extensively translocated throughout the plant, extremely stable in plant**
- **blocks synthesis of aromatic amino acids**
- **Very good for perennial species**
- **Weeds: Nonselective**
- **Used in renovation and dormant bermudagrass**
- **Dissipation via adsorption & microbial**



Roundup 32 oz, 19 DAT

3, 15, 23 Microtubule growth Inhibition

- plants grow by making new cells
 - process of cell division, mitosis
- plants are particularly susceptible as emerging seedlings
 - both shoot and roots
- newly forming roots can be susceptible at most stages of plant growth

Microtubule growth Inhibition

- most growth inhibition herbicides are soil applied and generally affect seedling weeds
- most interfere w/ mitosis (mitotic poisons)
- others appear to prevent lipid (cell membrane) production
- some prevent cell wall formation
- soil active, little movement once absorbed

Microtubule growth Inhibition


- *benefin* - PPI or PRE with irrigation
 - Balan
- *pendimethalin* – PRE
 - Prowl
- soil applied - annual grasses and certain broadleaf weeds
- vary in volatility and photodegradation
- prevent both root and shoot growth, inhibit cell division (mitosis)
- Very effective on small seeded weeds
- Plants cannot take up water-nutrients - starve

Microtubule growth Inhibition



4 – PGRs (plant growth regulators)

- *2,4-D, 2,4-DB, dicamba and more.....*
- BL weed control for a variety of crops (corn, pastures, legumes) and noncropland
- Cotton & tomato very sensitive – ppb range
- foliar & root uptake- extensive translocation
- interferes with nucleic acid (DNA and RNA) and protein synthesis; cells undergo rapid uncontrolled division and elongation



4- Benzoic Acids

- dicamba (broadleaf weeds)
- post-emergent soil active
- readily absorbed by leaf tips
- interferes with protein synthesis - leading to rapid, uncontrolled growth (similar to phenoxy)



Growth Regulator Herbicides

- **Phenoxy**
 - 2,4-D
- **Benzoics**
 - dicamba
- **Pyridines**
 - clopyralid
 - triclopyr
 - fluroxypyr



14 - PROTOX Inhibitors

- Biological activity
 - Mode of action - PPO or PROTOX inhibitors, contact action
 - require light for activity
 - Selectivity - metabolism



14 - PROTOX Inhibitors

- PRE & POST applied
- Depends on the chemistry
 - Flumioxazin (Chateau)
 - Carfentrazone (Aim)
- Flumioxazin - residual, rate dependent
- Carfentrazone no residual activity



fomesafen



Tomato



Ivyleaf morningglory

5, 6, 7, 22 Photosynthesis inhibitors



Metribuzin

Photosynthesis inhibitors

- **Biological activity**
 - Photosynthesis (PS I & II) inhibitors
 - readily absorbed by plant roots and translocated to leaves via transpiration stream
 - Selectivity based on metabolism
- **Dissipation**
 - Microbial
 - Hydrolysis
 - Soil & OM absorption
- **pH affects availability, increase pH, increase activity**
 - Metribuzin, WSSA Group 5

22- paraquat



Corn

Soybean



Herbicide chemistry

- **Water solubility is important**
 - Table salt 360 g/L = 3 lb/gal water
- **Glyphosate (K+ formulation)**
 - 10.5 g/L = 0.1 lb/gal water
- **Paraquat**
 - 620 g/L = 5.2 lb/gal water



Herbicide chemistry
pH and temperature effects

- **Water solubility is important**
 - Table salt 360 g/L = 3 lb/gal water
- **Metsulfuron – Patriot, multiple formulations**
 - pH 5.0 - 0.55 g/L = 0.0046 lb/gal water
 - pH 7.0 - 2.8 g/L = 0.023 lb/gal water
 - pH 9.0 - 213 g/L = 1.78 lb/gal water **350x**
- **Carfentrazone**
 - 68 F - 12 g/L = 0.1 lb/gal water
 - 86 F - 23 g/L = 0.2 lb/gal water



Herbicide chemistry

- **Water solubility is important**
 - Table salt 360 g/L = 3 lb/gal water
- **Flumioxazin –**
 - 0.00179 g/L = 0.000015 lb/gal
 - Low water solubility can lead to issues.....
 - Tank cleanout!!!




2019 Baleage and Silage Short Course:
The Uptake, Mode of Action, and Fate of
Herbicides in Hayfields

Dr. Tim Gray
UGA Prof. and Research Weed Scientist

Main points

- **Pesticides have to go somewhere!**
- Break down can be rapid in the environment
 - Depends on pesticide molecule chemistry:
 - Volatility
 - Solubility
 - Stability (resistance to photolysis, hydrolysis, etc.)
 - Depends on the environment (moisture, heat)
 - Depends on application method (granule, spray)
- Leaching
 - Need to move into treated soil
 - Do not want to move into ground water

Thank you



UNIVERSITY OF GEORGIA

Question????



UNIVERSITY OF GEORGIA
EXTENSION