

HAY MOISTURE: HOW DRY IS DRY ENOUGH?

June 2012 Hay & Forage Grower
Dennis Hancock, Forage Extension Specialist
The University of Georgia

There is a great misconception that once hay is “dry” and baled it is plain and devoid of life. The truth is that hay is never completely dry, and it is full of microscopic life. If the hay is not dry enough, those microscopic life forms can cause major problems.

Many microorganisms (mainly fungi species like *Aspergillus* and *Fusarium*, bacteria, and others) are ever present in hay (Figure 1). They feed on available carbohydrates on the surface of the forage plants and inside the stems and leaves. This feeding results in the loss of some dry matter (DM), reduces the quality of the hay, and also generates heat. The temperature of these hay bales, stacks, and barns can get very hot. In extreme cases, it can get so hot that the bales can spontaneously combust. Even if the temperature does not reach these extremes, these microorganisms can also form spores that give the hay a moldy smell and convert ordinary plant compounds into potent mycotoxins.




Figure 1. Summary of heating during hay storage, including recommended actions at various hay temperatures, what is causing the temperature increase, and what is happening as a result of the heat.

Nearly all hay goes through “a sweat” during the first few days after baling when the temperature rises. Figure 2 shows two cuttings of alfalfa hay for a study conducted at the University of Kentucky wherein the bales’ temperature was tracked over time. Notice that the

summer cutting, which was put up at 16% moisture, stayed relatively cool even during higher average ambient air temperatures. However, the fall cutting was baled a little wet (20% moisture) for round bales and it spiked over 140 °F within just 3 days.




Figure 2. Temperature of round bale alfalfa hay from summer (16% moisture) and fall (20% moisture) cuttings relative to the ambient air temperature during the first few days after baling.

The heat that is generated when hay goes through “a sweat” is a side effect of the microorganisms consuming the most digestible portions of the forage, such as carbohydrates like sugar and starch. Consequently, a substantial portion of the hay could be consumed and lost during this process.

Dr. Wayne Coblenz, Research Agronomist at the USDA-Agricultural Research Service’s U.S. Dairy Forage Research Center, has conducted several experiments on the impact that hay moisture and the resulting heating of the hay have on DM loss, hay quality, and heat risk. He recently found that for every 10°F increase in maximum temperature, the producer making alfalfa-orchardgrass hay would lose nearly 2% of the DM during storage.

It is worth noting, though, that some forage species have more readily available carbohydrates than others. As a result, some forage species may lose DM at a faster or slower rate. For example, Dr. Coblenz showed that the producer making bermudagrass hay would lose about 1.3% of the DM for every 10 °F increase in maximum temperature. This lower rate of DM loss is, in part, because bermudagrass has less readily-available carbohydrates.

Since these losses are coming from the most digestible forms of energy in the forage, hay heating comes at the expense of digestibility and the concentration of energy in the forage. Dr. Coblenz showed that the TDN of an alfalfa-orchardgrass hay lot is decreased by more than 2 percentage points and by more than 1 percentage points for bermudagrass hay for every 10 °F increase in maximum temperature. In other words, a good alfalfa-orchardgrass hay crop that was just a little too wet when it was baled might have went into the barn at 62% TDN, but it likely will come out of the barn with less than 56% TDN if it heated up to 140 °F or more.

In this article, the focus has been on limiting the maximum temperature, but it should be noted that the amount of time that the hay spends with an elevated temperature is also a concern. Like the chef who knows that whether by slow roasting for too long or cooking it too fast the meal can be ruined just the same, hay growers should minimize the time their hay spends at even moderate temperatures.

Much of the original research suggests hay moisture content should be kept less than 20% for small rectangular bales, less than 18% for round bales, and less than 16% for large rectangular bales. These are still good “rules of thumb,” but there are exceptions. Consider, for example, the advances in bale package sizes and high-density baling systems that have occurred in the modern era. In addition to the effects of bale type, size, and density, other factors contribute to the extent of hay heating. The amount of available carbohydrates in the forage crop, air circulation around/in the hay stack, relative humidity in the storage area, and the ambient temperature/humidity outside can each affect hay heating. Each producer’s situation will be somewhat different because of equipment, storage technique, and climatic differences. So, within the ranges provided in Figure 3, hay growers should allow for the effect that these factors might influence which target bale moisture is right for their farm.




Figure 3. The effect of bale moisture on the amount of damage that can be expected with different sizes and densities of hay bales, as well as other factors that affect hay heating.